



MLA AKT Practice Paper Exam Part 1

The MLA AKT Exam Board has put together a 200-item practice exam (2 x 100 item papers) to help medical students prepare for the UK Medical Licensing Assessment Applied Knowledge Test (MLA AKT). Blueprinted to the GMC Content Map this exam has been designed to reflect the style and type of question that students will encounter.

The practice exam comes with and without the answer options.

We hope students will find this a valuable resource.

Should you have any questions about the clinical content of the practice exam please speak to the Assessment Lead in your school in the first instance.

Any redistribution or reproduction of part or all of the contents in any form is prohibited other than for personal, educational and non-commercial use.



- 1.** A 79 year old woman has had a fall at home. She lives with her son and is dependent for shopping and finances. She has osteoarthritis and mild cognitive impairment.

She has bruises of different colours on her upper arms and torso. She avoids eye contact and becomes distressed when her son answers questions on her behalf. When spoken to alone, she says that she is 'a burden' and is frightened of being sent to a care home.

What is the most appropriate next step in management?

- A.** Arrange home visit follow up by GP
- B.** Discuss concerns with the son and seek his explanation
- C.** Make an adult safeguarding referral
- D.** Refer to falls prevention service
- E.** Request psychiatric assessment for capacity



2. A 29 year old man has 4 weeks of occasional painless rectal bleeding.

The perianal area and rectal examination are normal. Proctoscopy shows internal haemorrhoids, but no evidence of thrombosis.

Which is the most appropriate initial management?

- A.** Fibre supplementation
- B.** Rubber band ligation
- C.** Sclerotherapy
- D.** Topical corticosteroids
- E.** Topical nitrate



- 3.** An 80 year old woman has 18 months of a slowly progressive decline in short-term memory, with no clear onset. She often repeats herself and forgets conversations. She is less interested in going out and recently failed to recognise a nephew she had not seen in a few years. She has a history of mild hypertension and takes amlodipine and atorvastatin. Her daughter now visits her every day to help with cooking and medications.

Neurological examination is normal. Her Montreal cognitive assessment score is 22/30.

What is the most likely diagnosis?

- A.** Alzheimer dementia
- B.** Depression
- C.** Frontotemporal dementia
- D.** Mild cognitive impairment
- E.** Vascular dementia



4. A 45 year old man has exertional breathlessness and a daily cough productive of white sputum. He also has intermittent wheeze and chest tightness that vary from day to day and sometimes wake him at night. He has smoked 20 cigarettes daily since the age of 18.

He uses salbutamol several times daily despite satisfactory inhaler technique. He agrees to smoking cessation support.

Spirometry:

Pre bronchodilator		Post bronchodilator	
FEV	2.0 L (57% predicted)	2.5 L (71% predicted)	
FVC	4.0 L (80% predicted)	4.2 L (84% predicted)	
FEV ₁ /FVC	50%	60%	

Which is the most appropriate inhaled treatment to start?

- A. Long-acting β_2 -agonist with inhaled corticosteroid
- B. Long-acting β_2 -agonist with long-acting muscarinic antagonist
- C. Inhaled corticosteroid alone
- D. Long-acting β_2 -agonist alone
- E. Long-acting muscarinic antagonist alone



- 5.** A 38 year old woman has 2 days of right upper quadrant pain, vomiting and fever. She has had intermittent pain for the past 3 months, worse after eating. Her BMI is 34 kg/m².

Her temperature 38.1°C, pulse 102 bpm, BP 100/72 mmHg and oxygen saturation 98% breathing air. She has right upper quadrant tenderness, but no hepatomegaly, and her abdomen is soft with normal bowel sounds.

Investigations:

Alanine aminotransferase (ALT) 42 IU/L (10–50)

Albumin 37 g/L (35–50)

Alkaline phosphatase 165 IU/L (25–115)

Aspartate aminotransferase (AST) 35 IU/L (10–40)

Bilirubin (total) 21 µmol/L (< 21)

Amylase 192 U/L (<220)

What is the most likely diagnosis?

- A.** Acute pancreatitis
- B.** Biliary colic
- C.** Cholecystitis
- D.** Duodenal ulcer
- E.** Hepatitis



- 6.** A 28 year old woman has 4 weeks of retrosternal pain when swallowing liquids or solids. She has asthma, treated with inhaled beclometasone with formoterol.

Investigations:

Gastroscopy: patchy white exudate scattered along length of oesophagus

What is the most likely type of oesophagitis?

- A.** Candida oesophagitis
- B.** Cytomegalovirus oesophagitis
- C.** Eosinophilic oesophagitis
- D.** Herpes simplex oesophagitis
- E.** Reflux oesophagitis



7. An 84 year old man presents to the ambulatory care unit following a fall. He reports several falls over the past 2 months, usually after getting out of bed to pass urine. He feels light-headed on standing, without loss of consciousness.

He has hypertension, atrial fibrillation and benign prostatic hyperplasia. He takes bisoprolol fumarate, finasteride, indapamide, tamsulosin hydrochloride and apixaban. Tamsulosin was added 2 months ago because of persistent urinary symptoms. His other medications have been unchanged for over a year.

His pulse is 76 bpm and irregular. His BP is 138/84 mmHg lying and 114/62 mmHg after standing for 1 minute, reproducing his light-headedness.

Investigations:

Haemoglobin 124 g/L (130–175)

White cell count $5.6 \times 10^9/L$ (3.8–10.0)

Platelets $164 \times 10^9/L$ (150–400)

Which medication is most likely to be contributing to his falls?

- A. Apixaban
- B. Bisoprolol
- C. Finasteride
- D. Indapamide
- E. Tamsulosin



- 8.** A 74 year old woman briefly loses consciousness whilst walking uphill. She recovers promptly without confusion. She has had chest discomfort and increasing breathlessness on exertion for 6 months.

Her pulse is 78 bpm and low volume, and BP 120/95 mmHg. Her chest is clear. She has an ejection systolic murmur that radiates to the carotids.

Investigations:

ECG: sinus rhythm with voltage criteria for left ventricular hypertrophy.

Which feature of her collapse is a red flag for a potentially serious cardiac cause?

- A.** Associated pallor
- B.** Brief duration of unconsciousness
- C.** Occurrence during exertion
- D.** Rapid spontaneous recovery
- E.** Recovery without confusion



- 9.** An 89 year old woman is breathless. Her medication includes furosemide 40 mg, ramipril 1.25 mg and bisoprolol fumarate 1.25 mg.

She has pitting oedema up to her thighs. Her JVP is elevated at 8 cm. She has fine inspiratory crepitations up to the mid-zones bilaterally.

Investigations:

Sodium 128 mmol/L (135–146)

Potassium 4.1 mmol/L (3.5–5.3)

Urea 5.6 mmol/L (2.5–7.8)

Creatinine 75 μ mol/L (60–120)

What is the most appropriate next step?

- A.** Give intravenous 0.9% sodium chloride
- B.** Give intravenous 1.8% sodium chloride
- C.** Increase furosemide dose
- D.** Increase ramipril dose
- E.** Stop bisoprolol fumarate



- 10.** An 82 year old man has had three episodes of brief loss of consciousness in the last 24 hours. None of these were associated with convulsions. He is now alert and does not recall any weakness or loss of sensation during the episodes. He is a nursing home resident and his carer reports that he became pale before losing consciousness and recovered quickly within seconds with flushing of his face. He has a history of dementia and hypertension.

His pulse rate during the episodes was 40 bpm but is now 80 bpm. His heart sounds are normal. He has normal strength, sensation, reflexes and visual fields.

Investigations:

ECG: Sinus rhythm, rate 74 bpm.

Which investigation is most likely to confirm the diagnosis?

- A.** 48 hour ECG monitoring
- B.** Carotid Doppler ultrasound
- C.** CT scan of head
- D.** EEG
- E.** Tilt table test



- 11.** A 74 year old man has had 2 hours of chest pain. He has hypertension and hypercholesterolaemia, and takes ramipril and atorvastatin. He smokes 30 cigarettes per day.

His pulse is 75 bpm and regular, BP 110/60 mmHg and oxygen saturation is 96% breathing air.

Investigations:

ECG: see image

Primary percutaneous coronary intervention is planned.



Which coronary artery is most likely to be occluded?

- A.** Circumflex
- B.** Left anterior descending
- C.** Left main
- D.** Obtuse marginal
- E.** Right



- 12.** A 34 year old woman collapses at the restaurant she works in. She told a colleague that she did not feel well and then dropped to the floor. Her colleagues report that she went pale and then her limbs jerked briefly when she was on the floor. She regained consciousness after 15 seconds and felt nauseous. She was not confused.

When paramedics arrive, her pulse is 89 bpm, BP 115/60 mmHg and oxygen saturation 98% breathing air.

Which is the most likely diagnosis?

- A.** Carotid sinus syncope
- B.** Focal seizure
- C.** Postural tachycardia syndrome
- D.** TIA
- E.** Vasovagal syncope



- 13.** A 90 year old man attends hospital following a fall. He has been struggling at home with washing, dressing and meal preparation, and now needs a package of care.

Which is the best tool to assess his activities of daily living?

- A.** Barthel Index
- B.** Malnutrition Universal Screening Tool
- C.** Montreal Cognitive Assessment
- D.** Rockwood Clinical Frailty Scale
- E.** Waterlow Score



- 14.** A 72 year old woman has 5 weeks of diarrhoea. She has fresh blood mixed with her stool and a sense of incomplete emptying of her rectum. She has no pain and has weight loss of 3 kg over 2 months.

Which is the most likely diagnosis?

- A.** Anal fissure
- B.** Colorectal cancer
- C.** Crohn's disease
- D.** Diverticulitis
- E.** Ulcerative colitis



- 15.** A 53 year old man has severe upper abdominal pain. He has loose, pale diarrhoea, and is passing bulky stools that flush away poorly and float in the toilet. He has had recurrent admissions with acute abdominal pain over the last 15 years. He has a long history of excess alcohol intake.

Investigation:

CT scan abdomen: calcification in the epigastric region.

Which is the most likely mechanism for his diarrhoea?

- A.** Bacterial overgrowth
- B.** Failure of duodenal cholecystokinin
- C.** Increased small bowel motility
- D.** Loss of pancreatic exocrine function
- E.** Reduced bile production



- 16.** A 77 year old man is admitted with 1 day of severe left iliac fossa pain and fresh rectal bleeding.

His temperature is 38.7°C, pulse 109 bpm and regular, and BP 119/64 mmHg. There is tenderness in the left iliac fossa, without guarding or rebound tenderness.

What investigation is the most appropriate to establish the underlying diagnosis?

- A.** Blood cultures
- B.** Contrast-enhanced CT scan of abdomen and pelvis
- C.** Flexible sigmoidoscopy
- D.** Stool culture
- E.** Ultrasound scan of abdomen



- 17.** A 59 year old woman has had 6 months of tiredness and aching in both legs, worse on the left than right. The symptoms are worse when she has been standing, and are relieved by elevating her legs. She has a history of COPD.

Her BP is 144/86 mmHg and JVP 2 cm above the sternal angle. She has telangiectasia and superficial varicosities on both lower legs with pitting oedema at the ankle. She has a 1.5 cm superficial, slightly tender ulcer behind the left medial malleolus. Her femoral and popliteal pulses are palpable. Her foot pulses are weakly palpable.

Which is the most likely diagnosis?

- A.** Bowen's disease
- B.** Chronic venous insufficiency
- C.** Lymphoedema
- D.** Peripheral arterial disease
- E.** Right heart failure



- 18.** A 25 year old man has a cough that has been keeping him awake at night for 8 weeks. He has recently had a cold that has now resolved. He has had a similar cough when exercising, which is worse in the winter. He has never smoked.

Which is the most likely diagnosis?

- A.** Asthma
- B.** Bronchiectasis
- C.** Chronic obstructive pulmonary disease
- D.** Sarcoidosis
- E.** Viral induced wheeze



- 19.** A 62 year old woman has 3 months of cough productive of green sputum. She has no fever, sweats or haemoptysis. She had pneumonia as a child. She has never smoked.

There are coarse crackles over both lungs. She has bilateral leg oedema. Sputum culture grows *Pseudomonas aeruginosa*.

What is the most likely diagnosis?

- A.** Bronchiectasis
- B.** Congestive cardiac failure
- C.** COPD
- D.** Lung cancer
- E.** Tuberculosis



- 20.** A 24 year old man has sharp stabbing right sided chest pain, worse when he breathes in. It began suddenly 2 hours ago with no history of trauma, but the pain has been worsening and he has noticed that he is also becoming breathless. He has a history of asthma and uses a salbutamol inhaler as required.

His pulse is 98 bpm, BP 128/89 mmHg, respiratory rate 24 breaths per minute and oxygen saturation 88% in air. There are decreased breath sounds on the right and the trachea is central.

Which is the most appropriate next step?

- A.** 15 L/min oxygen by reservoir bag mask
- B.** 24% oxygen by Venturi mask
- C.** Arterial blood gases
- D.** Right sided intercostal drain
- E.** Right sided needle thoracocentesis



- 21.** A 68 year old woman has 6 weeks of cough. She also has fatigue and has lost 4 kg in weight over 3 months. She is a smoker with a 30 pack-year history.

She has no palpable lymph nodes. Chest and abdominal examinations are unremarkable.

Investigations:

Haemoglobin 120 g/L (115–150)

White cell count $6.4 \times 10^9/L$ (3.8–10.0)

Platelets $290 \times 10^9/L$ (150–400)

Chest X-ray: 2-cm mass in upper lobe of right lung

Which is the most appropriate next investigation?

- A.** Bronchoscopy
- B.** CT of chest and abdomen
- C.** Isotope bone scan
- D.** PET-CT
- E.** Pulmonary function tests



- 22.** A 65 year old woman has 1 day of breathlessness, 1 week of progressive headache and 3 months of weight loss and cough.

Her pulse rate is 88 bpm, respiratory rate 20 breaths per minute and oxygen saturation 95% breathing oxygen 4L/min via nasal prongs. She has a swollen face and neck and distended veins on her chest. There are reduced breath sounds at the right upper lobe. Chest X-ray shows a mass in the right upper lobe.

Which is the most appropriate diagnostic investigation?

- A.** Carotid Doppler scan
- B.** Contrast-enhanced CT scan of chest
- C.** CT scan of head
- D.** Duplex ultrasound scan of subclavian vein
- E.** Venography



- 23.** A 44 year old man has just finished treatment with IV antibiotics for an episode of spontaneous bacterial peritonitis. He has alcohol related liver disease and is being considered for a liver transplant. His medications include spironolactone, furosemide and thiamine.

His temperature is 36.4°C, pulse 76 bpm and BP 104/64 mmHg. He has shifting dullness.

Which additional treatment should be started?

- A.** Bendroflumethiazide
- B.** Ciprofloxacin
- C.** Midodrine hydrochloride
- D.** Prednisolone
- E.** Propranolol hydrochloride



24. An 84 year old woman has had 8 months of malaise and lethargy.

She is pale. Her pulse is 86 bpm and BP 144/94 mmHg. Her chest is clear.

Investigations:

Haemoglobin 90 g/L (115–150)

Mean cell volume (MCV) 84 fL (80–96)

Urea 15.7 mmol/L (2.5–7.8)

Creatinine 260 $\mu\text{mol/L}$ (60–120)

eGFR 16 mL/min/1.73 m² (>60)

Serum vitamin B₁₂ 180 ng/L (160–925)

Serum folate 8 $\mu\text{g/L}$ (3–15)

Ferritin 10 $\mu\text{g/L}$ (12–200)

Which is the most appropriate initial treatment for her anaemia?

- A.** Blood transfusion
- B.** Dialysis
- C.** Erythropoietin
- D.** Iron replacement
- E.** Thiamine



- 25.** A 74 year old woman attends her GP with jaundice, generalised itching and dark urine for 4 days. She completed a 10-day course of co-amoxiclav for pneumonia 1 week ago.

Investigations:

Albumin 41 g/L (35–50)

Total bilirubin 150 $\mu\text{mol/L}$ (<21)

Alanine aminotransferase 101 IU/L (10–50)

Alkaline phosphatase 600 IU/L (25–115)

Ultrasound scan: no gallstones or biliary duct dilatation.

What is the most likely cause of her jaundice?

- A.** Alcoholic-related hepatitis
- B.** Choledocholithiasis
- C.** Drug-induced cholestatic hepatitis
- D.** Liver abscess
- E.** Primary biliary cholangitis



- 26.** A 56 year old woman has had 6 months of malaise and night sweats. She does not smoke and drinks 20 units of alcohol per week.

She has jaundice.

Investigations:

AST 133 IU/L (10–40)

ALP 114 IU/L (25–115)

Bilirubin 45 μ mol/L (<17)

γ GT 150 IU/L (9–40)

Antinuclear antibodies (ANA) 1:640 (<20)

Antimitochondrial antibodies negative

Anti-smooth muscle antibodies positive

What is the most likely diagnosis?

- A.** Alcoholic hepatitis
- B.** Autoimmune hepatitis
- C.** Hepatitis C infection
- D.** Primary biliary cholangitis
- E.** Primary sclerosing cholangitis



27. A 32 year old man is referred for genetic counselling following a recent myocardial infarction. At the time of admission he was noted to have a total cholesterol of 9.6 mmol/L (<5.0) and LDL cholesterol of 5 mmol/L (<3.0). His uncle died of a heart attack aged 41 years and his grandmother died of a stroke aged 46 years. He is afraid that he may pass on his risk of heart disease to any future children.

Which is the most likely pattern of inheritance for this condition?

- A.** Autosomal dominant
- B.** Autosomal recessive
- C.** Mitochondrial
- D.** X-linked dominant
- E.** X-linked recessive



- 28.** A 60 year old man collapses in a shopping centre and bystanders initiate basic life support. Upon arrival, the paramedics find the patient unresponsive, without a pulse. A cardiac monitor is attached and rhythm strip obtained (see image).



Which is the most appropriate initial intervention?

- A.** Defibrillation
- B.** Intravenous adenosine
- C.** Intravenous amiodarone hydrochloride
- D.** Intravenous epinephrine
- E.** Synchronised cardioversion



- 29.** An 81 year old woman has had 6 months of intermittent visual hallucinations where she sees small children and farmyard animals. They are more often seen at home and in the evenings. She does not recognise any of the children and they do not interact with her. She describes no auditory, olfactory or tactile hallucinations, and she feels that her thoughts are her own. She is otherwise well and is not concerned about her mood or memory. She is registered partially sighted because of macular degeneration. She has a history of hypertension.

Her visual acuity is 6/36 in both eyes. She has no other focal neurological deficit. Her abbreviated mental test score is 10/10. CT scan of head shows small vessel disease only.

Which is the most likely diagnosis?

- A.** Charles Bonnet syndrome
- B.** Dementia with Lewy bodies
- C.** Focal aware seizures
- D.** Occipital stroke
- E.** Schizophrenia



- 30.** A 70 year old man has 2 days of confusion, disorientation, difficulty with balance and visual disturbance, of sudden onset. He has AF and hypertension.

His temperature is 37.2°C, pulse 98 bpm, BP 128/70 mmHg, respiratory rate 15 breaths per minute and oxygen saturation 96% breathing air.

Investigations:

CT scan of head: see image



What is the most likely diagnosis?

- A.** Left cerebellar infarction
- B.** Left occipital infarction
- C.** Left occipital metastasis
- D.** Left parietal abscess
- E.** Left parietal infarction



31. A 75 year old man is admitted to hospital with pneumonia and reduction in his mobility. He takes amlodipine, apixaban, atorvastatin, bisoprolol fumarate, metformin hydrochloride and ramipril.

His eGFR is 37 mL/min/1.73 m² (>60). His routine medication and an antibiotic have been prescribed.

Which is the most appropriate management to reduce the risk of venous thromboembolism?

- A.** Aspirin
- B.** Low molecular weight heparin
- C.** No additional treatment
- D.** Tranexamic acid
- E.** Unfractionated heparin



- 32.** A 17 year old girl has 2 days of loss of appetite, nausea and central to lower abdominal pain. Her last menstrual period was 30 days ago.

Her temperature is 37.7°C. She is tender in the right Iliac fossa and hypogastrium. There is no lymphadenopathy.

Investigations:

White cell count $12.3 \times 10^9/L$ (3.8–10.0)

Pregnancy test: negative

Urinalysis: blood negative, protein negative

Which is the most likely diagnosis?

- A.** Acute appendicitis
- B.** Acute cholecystitis
- C.** Ectopic pregnancy
- D.** Inflammatory bowel disease
- E.** Pelvic Inflammatory disease



- 33.** A 59 year old man has 6 months of bilateral calf pain on walking around 100 metres. He has CKD stage 5, hypertension and type 2 diabetes mellitus. His medication includes amlodipine, doxazosin and ramipril.

His BP is 142/90 mmHg. His femoral pulses are palpable. His popliteal and foot pulses are difficult to feel. He has pitting oedema.

Investigations:

Ankle brachial pressure index (ABPI): right leg 1.4, left leg 1.4

What is the most likely cause of his ABPI result?

- A.** Capillary advanced glycation end production
- B.** Incompetent venous valves
- C.** Increased aortic elasticity
- D.** Increased interstitial hydrostatic pressure
- E.** Medial arterial calcification



- 34.** A 56 year old man has had fever, malaise and increasing breathlessness for 2 weeks. He has been off work with a cough and general ill health for three weeks. He has a 25 pack year smoking history.

He has a temperature of 38.3°C, pulse 98 bpm, BP 106/78 mmHg and respiratory rate of 24 breaths per minute. His oxygen saturation is 94% breathing air. There is reduced movement of the right chest with dullness to percussion, reduced breath sounds and absent vocal resonance over the right lung base.

Investigations:

White cell count $26 \times 10^9/L$ (3.0–10.0)

Serum C reactive protein 350 mg/L (<5)

What is the most likely diagnosis?

- A.** Bronchial carcinoma
- B.** Empyema
- C.** Infective exacerbation of COPD
- D.** Lobar pneumonia
- E.** Pneumothorax



- 35.** A 42 year old man has 3 days of increasing breathlessness and a cough productive of blood-streaked, green sputum.

His temperature is 39.2°C, pulse 105 bpm, BP 105/60 mmHg, respiratory rate 24 breaths per minute and oxygen saturation 91% breathing air. There is reduced air entry with coarse crackles at the left lung base. The skin of the palms of his hands is peeling.

Chest X-ray: bilateral consolidation with multiple cavities

Which is the most likely causative organism?

- A.** *Haemophilus influenzae*
- B.** *Legionella pneumophila*
- C.** *Mycobacterium tuberculosis*
- D.** *Staphylococcus aureus*
- E.** *Streptococcus pneumoniae*



- 36.** A 77 year old man has had worsening breathlessness on exertion, cough and tiredness for 6 months. He is a retired plumber with a 50 pack-year history of cigarette smoking.

He is cachectic. His temperature is 36.6° C, pulse 94 bpm, BP 100/72 mmHg, respiratory rate 16 breaths per minute and oxygen saturation is 89% breathing air. He has reduced expansion and quiet breath sounds on auscultation of the left hemithorax.

Investigations:

Chest X-ray: diaphragmatic pleural plaques. Lobulated thickening on the left pleura with small pleural effusion. Reduced volume of the left hemithorax.

What is the most likely diagnosis?

- A.** Bronchial carcinoma
- B.** Mesothelioma
- C.** Parapneumonic effusion
- D.** Pleural metastases
- E.** Pneumothorax



37. A 47 year old man has had 3 months of dull, intermittent epigastric pain. It is worse in the morning and relieved by drinking milk. His weight is steady. He drinks 20 units of alcohol per week.

His abdomen is soft, with some discomfort in the epigastrium on deep palpation. There are no masses.

Investigations:

Haemoglobin 127 g/L (130–175)

MCV 77 fL (80–96)

Which is the most likely diagnosis?

- A.** Chronic pancreatitis
- B.** Coeliac disease
- C.** Duodenal ulcer
- D.** Gastro-oesophageal reflux
- E.** Pancreatic cancer



- 38.** A 60 year old woman develops wheeze and chest tightness after discharge from hospital following a myocardial infarction. She has COPD, type 2 diabetes mellitus and depression. Her medications include aspirin, atorvastatin, bisoprolol, clopidogrel, ramipril and sertraline.

Her pulse is 64 bpm, BP 124/76 mmHg and oxygen saturation 96% breathing air. There is widespread expiratory wheeze without crackles. Her JVP is 2 cm above the sternal angle.

Which medication is most likely to have caused her new respiratory symptoms?

- A.** Atorvastatin
- B.** Bisoprolol
- C.** Clopidogrel
- D.** Ramipril
- E.** Sertraline



- 39.** A 19 year old woman has had severe abdominal pain and passing bloody diarrhoea 6 to 8 times per day for the past 3 days. She has ulcerative colitis and takes sulfasalazine.

Her temperature is 37.9°C, pulse 100 bpm and regular, BP 110/60 mmHg and oxygen saturation 98% breathing air. Her abdomen is soft and mildly tender throughout. There is no distension, guarding or rebound tenderness, or organomegaly. She has active bowel sounds.

Intravenous fluids have been started.

Which is the most appropriate initial treatment?

- A.** Intravenous ciclosporin
- B.** Intravenous hydrocortisone
- C.** Intravenous infliximab
- D.** Oral loperamide hydrochloride
- E.** Rectal mesalazine



- 40.** A 25 year old woman has had recurrent episodes of abdominal discomfort associated with loose and urgent bowel movements for a year. There is no blood in the stool and her weight is stable.

Her inflammatory markers, coeliac screen and faecal calprotectin are all normal. Faecal immunochemical test negative.

She is given lifestyle and dietary advice.

What is the most appropriate next step in management?

- A.** Advise her to buy ispaghula husk over the counter
- B.** Prescribe amitriptyline
- C.** Prescribe fluoxetine
- D.** Prescribe loperamide
- E.** Refer for a colonoscopy



- 41.** A 25 year old student has had urgency and loose stools for several days. Several of her friends have had a similar illness. It is the end of term and she needs to make a 5 hour car journey home.

What is the most appropriate medication?

- A.** Lactulose
- B.** Loperamide
- C.** Mebeverine
- D.** Mesalazine
- E.** Prochlorperazine



- 42.** A 54 year old man has had worsening shortness of breath and swollen legs for 2 months. He has severe pulmonary fibrosis.

His temperature is 37.1°C, pulse 100 bpm, BP 114/86 mmHg, respiratory rate 17 breaths per minute and oxygen saturation 90% breathing air. There is a pansystolic murmur loudest in inspiration at the left sternal edge. There are fine inspiratory crackles throughout the chest. There is swelling of both legs up to the calf.

What is the most likely diagnosis?

- A.** Congestive cardiac failure
- B.** Infective endocarditis
- C.** Pulmonary embolus
- D.** Pulmonary hypertension
- E.** Rheumatic mitral valve disease



- 43.** A 74 year old man is admitted for investigation of chest pain. He has a history of dementia. All investigations are normal, but at night he is wandering around the ward and refusing to return to his bed.

What is the most appropriate first line treatment?

- A.** Donepezil
- B.** Haloperidol
- C.** Lorazepam
- D.** Orientation strategies
- E.** Zopiclone at night



44. An 84 year old man has agitation and is unable to weight bear, two days after admission following a fall at home. When assessed by the physiotherapist, he is unwilling to stand up and shouts loudly when his right leg is touched. He is prescribed regular analgesia.

Admission X-rays of his hips and pelvis show only mild degenerative changes.

Which is the most appropriate next action?

- A.** CT scan of hips and pelvis
- B.** Encourage mobilisation
- C.** Lorazepam as required
- D.** Referral to psychiatry of old age
- E.** Repeat hip and pelvic X-ray



- 45.** A 51 year old man develops a distended abdomen 4 days after surgery for a perforated appendix. He has not passed flatus for 24 hours. He is being treated with oral paracetamol, and intravenous ceftriaxone and metronidazole.

His temperature is 37.4°C. His abdomen is soft and non-tender but distended, with scanty bowel sounds.

What is the most likely diagnosis?

- A.** Pelvic collection
- B.** Postoperative ileus
- C.** Pseudomembranous colitis
- D.** Small bowel ischaemia
- E.** Small bowel obstruction



46. A 40 year old man has become gradually more unwell over 2 days. He feels very tired, weak and short of breath. He does not have a cough. He has schizophrenia and started treatment with clozapine 3 weeks ago. He has no history of medical problems and is a non-smoker.

His temperature is 37.5°C, pulse 105 bpm, BP 120/80 mmHg and respiratory rate is 25 breaths per minute. His heart sounds are normal and his chest is clear.

Investigations:

Haemoglobin 140 g/L (130–175)

White cell count $12.1 \times 10^9/L$ (3.8–10.0)

Creatine kinase 400 U/L (25–200)

Troponin I 2168 ng/L (< 5)

ECG: widespread concave ST segment elevation with T-wave inversion

Chest X-ray: no abnormalities

What is the most likely diagnosis?

- A.** COVID-19 pneumonia
- B.** Myocardial infarction
- C.** Myocarditis
- D.** Neuroleptic malignant syndrome
- E.** Sepsis



- 47.** A 54 year old man has worsening jaundice, nausea and fever for 3 days. He has had a lack of appetite for 3 weeks. He drinks 50 units of alcohol a week and smokes 10 cigarettes a day. He has hypertension and hypercholesterolaemia. He takes amlodipine and atorvastatin.

His temperature is 38.0°C, pulse 100 bpm, BP 100/60 mmHg, respiratory rate 16 breaths per minute and oxygen saturation is 100% breathing air. His BMI is 25 kg/m². He is jaundiced and has spider naevi. His abdomen is soft and non-tender.

Investigations:

Alanine aminotransferase (ALT) 80 IU/L (10–50)

Albumin 32 g/L (35–50)

Alkaline phosphatase (ALP) 170 IU/L (25–115)

Aspartate aminotransferase (AST) 180 IU/L (10–40)

Bilirubin (total) 90 µmol/L (<21)

Gamma glutamyl transferase (γGT) 220 IU/L (9–40)

What is the most likely cause for his current presentation?

- A.** Acute viral hepatitis
- B.** Alcoholic hepatitis
- C.** Autoimmune hepatitis
- D.** Drug-induced hepatitis
- E.** Metabolic dysfunction-associated steatotic liver disease



- 48.** A 93 year old woman has reduced consciousness and difficulty swallowing. She has developed aspiration pneumonia and delirium. She has hypertension and dementia. She usually takes amlodipine, donepezil and ramipril.

Her temperature is 38.0°C, pulse 114 bpm, BP 107/76 mmHg, respiratory rate 18 breaths per minute and oxygen saturation 91% breathing air.

Investigations:

Sodium 151 mmol/L (135–146)

Potassium 5.4 mmol/L (3.5–5.3)

Urea 19.6 mmol/L (2.5–7.8)

Creatinine 135 µmol/L (60–120)

eGFR (now) 35 mL/min/1.73 m² (>60)

eGFR (1 month ago) 42 mL/min/1.73 m² (>60)

Urinary osmolality 750 mosmol/kg (350–1000)

What is the most likely cause of her hypernatraemia?

- A.** Acute kidney injury
- B.** Adrenal insufficiency
- C.** Dehydration
- D.** Ramipril
- E.** Syndrome of inappropriate antidiuresis



- 49.** A 35 year old woman has sudden onset of breathlessness and right sided pleuritic chest pain. She was recently found to have bony metastases from previously treated breast carcinoma and received her second cycle of chemotherapy 2 days ago. She has a history of asthma.

Her temperature is 37.3°C, pulse 102 bpm, BP 105/60 mmHg, respiratory rate 26 breaths per minute and oxygen saturation 94% in air. Her chest is clear.

What is the most likely diagnosis?

- A.** Lung metastases
- B.** Lymphangitis carcinomatosa
- C.** Pneumonia
- D.** Pneumothorax
- E.** Pulmonary embolus



50. A 61 year old man is in atrial fibrillation. He was previously well.

His pulse rate is 76 bpm and BP 121/74 mmHg. An echocardiogram is normal. He does not want cardioversion.

Which is the most appropriate management option?

- A.** Apixaban
- B.** Aspirin
- C.** Clopidogrel
- D.** No medication required
- E.** Warfarin sodium



51. A 64 year old man with type 2 diabetes mellitus attends for review. He takes metformin 1000 mg daily and the maximum tolerated dose of losartan.

His BP is 128/76 mmHg and BMI is 30 kg/m². He has no peripheral oedema.

Investigations:

HbA1c 49 mmol/mol (20–42)

Creatinine 128 µmol/L (60–110) (creatinine 122 µmol/L 6 months ago)

Estimated GFR 51 mL/min/1.73 m² (>60)

Potassium 4.5 mmol/L (3.5–4.9)

Urine albumin ratio 38 mg/mmol (<3)

Which treatment change is most appropriate to reduce his risk of chronic kidney disease progression?

- A.** Add a DPP-4 inhibitor
- B.** Add an SGLT2 inhibitor
- C.** Add a thiazide-like diuretic
- D.** Increase the dose of metformin
- E.** Switch losartan to ACE inhibitor



52. A 34 year old man has fever and a cough productive of brown sputum, flecked with blood for one week. He has right sided pleuritic chest pain. He smokes and injects heroin.

He has coarse crackles in his right lung and a sinus over his right femoral vein.

Investigation:

CT Pulmonary Angiogram shows right sided consolidation and some early cavitation with right sided pulmonary emboli.

What is the most likely diagnosis?

- A.** Diamorphine induced chronic pulmonary disease
- B.** Inferior vena cava thrombosis
- C.** Pulmonary haemorrhage
- D.** Pulmonary tuberculosis
- E.** Tricuspid endocarditis



- 53.** A 48 year old woman is reviewed in the renal clinic. She has CKD stage 4. Her medications include colecalciferol.

Investigations:

Urea 17.3 mmol/L (2.5–7.8)

Creatinine 295 μ mol/L (60–120)

eGFR 17 mL/min/1.73 m² (>60)

ALP 120 IU/L (25–115)

Calcium (adjusted) 2.3 mmol/L (2.2–2.6)

Phosphate 2.8 mmol/L (0.8–1.5)

Vitamin D 78 nmol/L (75–100)

Parathyroid hormone 60.0 pmol/L (1.6–8.5)

Which is the most appropriate class of treatment to start?

- A.** Bisphosphonate
- B.** Calcimimetic
- C.** Calcitonin
- D.** Phosphate binder
- E.** Vitamin D analogue



- 54.** A 24 year old man has had flu-like symptoms for 4 weeks. He has been taking paracetamol twice a day for his fevers, which lasted 2 weeks and stopped 10 days ago. He drinks 12 units of alcohol a week.

His temperature is 37.0°C, pulse 80 bpm and BP 120/70 mmHg. His abdomen is normal.

Investigations:

Haemoglobin 140 g/L (130–175)

White cell count $3.8 \times 10^9/L$ (4.0–11.0)

Platelets $200 \times 10^9/L$ (150–400)

Alanine aminotransferase (ALT) 186 IU/L (10–50)

Alkaline phosphatase (ALP) 75 IU/L (25–115)

Bilirubin (total) 30 $\mu\text{mol/L}$ (<21)

Which is the most likely underlying cause?

- A.** Alcoholic hepatitis
- B.** Drug-induced hepatitis
- C.** Epstein Barr virus
- D.** Gilbert's syndrome
- E.** Hepatitis A



- 55.** A 25 year old man has blood borne virus screening before starting medical school. Nine months previously he had fatigue, nausea and poor appetite for 2 weeks. Investigations at that time showed a positive hepatitis B surface antigen result. He currently feels well.

Investigations:

Hepatitis B surface antigen (HBsAg): positive

Hepatitis B e antigen (HBeAg): positive

Antibody to hepatitis B surface antigen (anti-HBs): positive

Antibody to hepatitis B core antigen (anti-HBc): IgG positive, IgM negative

Antibody to hepatitis B e antigen (anti-HBe): negative

Hepatitis B virus DNA: detectable

Which is the best explanation of these results?

- A.** Acquired immunity to hepatitis B infection
- B.** Acute hepatitis B infection
- C.** Chronic hepatitis B infection
- D.** Inactive hepatitis B infection
- E.** Previous vaccination against hepatitis B



56. A 74 year old man becomes very sweaty and then loses consciousness on a postoperative surgical ward. He has a history of type 2 diabetes treated with insulin. His immediate capillary blood glucose is 2.2 mmol/L, and he is given 100 mL of 20% dextrose.

He is now conscious and talking coherently. His repeat capillary blood glucose is 4.9 mmol/L.

Which is the most appropriate next step in management?

- A.** Administer 1 L of 5% dextrose over 8 h
- B.** Administer intramuscular glucagon 1 mg
- C.** Give him a carbohydrate snack
- D.** Give him a cup of tea with two sugars
- E.** Omit the next dose of insulin



57. A 51 year old woman has had dizziness and unsteadiness for 1 month. Her symptoms are present most of the time and there are no clear precipitants.

Her BP is 146/87 mmHg. She has reduced hearing in the right ear. She has nystagmus on right lateral gaze. She has no diplopia.

Which is the most appropriate management?

- A.** CT scan of head
- B.** MR scan of brain
- C.** Perform Dix–Hallpike manoeuvre
- D.** Start aspirin
- E.** Start betahistine



58. A 32 year old man attends his GP with nausea, vomiting and worsening abdominal pain which started 12 hours ago. The pain is on the left side of his abdomen which is not relieved by paracetamol. He has dysuria and the urine appears dark and cloudy. He has rheumatoid arthritis and takes oral methotrexate weekly.

His temperature is 38.2°C, pulse 114 bpm, BP 108/58 mmHg and oxygen saturation 98% breathing air. His abdomen is soft with tenderness over the left flank.

Urinalysis: blood 2+, protein 2+, leucocytes 2+

Which is the most appropriate next step in management?

- A.** Blood and urine cultures
- B.** Oral co-amoxiclav
- C.** Oral nitrofurantoin
- D.** Urgent hospital admission
- E.** Urgent US scan of abdomen



59. A 65 year old white woman has itching on both lower legs. She is under the palliative care team with ovarian cancer with metastases but is not acutely unwell.

She has bilateral oedema to the knees. The skin on the legs is dry, red and scaly.

Which is the most likely diagnosis?

- A.** Bilateral cellulitis
- B.** Erythema nodosum
- C.** Lymphoedema from pelvic compression
- D.** Paraneoplastic deep vein thromboses
- E.** Venous eczema



60. An 80 year old man has multiple lumps on his bald scalp.

There are numerous erythematous keratotic lesions that are very rough on palpation. The surrounding skin is freckled, with focal areas of hypopigmentation.

Which is the most likely neoplastic complication of this condition?

- A.** Basal cell carcinoma
- B.** Kaposi's sarcoma
- C.** Malignant melanoma
- D.** Mycosis fungoides
- E.** Squamous cell carcinoma



61. A 62 year old woman has a rash on her neck and upper chest which is made worse by sun exposure. She also has fatigue and difficulty climbing the stairs at home. She has no significant past medical history and takes no regular medication. She has a symmetrical purplish discoloration of her eyelids with mild eyelid oedema. She has grade 4/5 proximal muscle weakness.

Which is the most likely diagnosis?

- A.** Atopic eczema
- B.** Dermatomyositis
- C.** Psoriasis
- D.** Rosacea
- E.** Systemic lupus erythematosus



62. A 52 year old man visits his GP with 2 months of general lethargy, weakness and leg cramps. He has a history of hypertension, heart failure and Barrett's oesophagitis. He takes amlodipine, bisoprolol, eplerenone, esomeprazole and losartan.

Investigations:

Sodium 139 mmol/L (135–146)

Potassium 4.3 mmol/L (3.5–5.3)

Calcium 2.2 mmol/L (2.2–2.6)

Magnesium 0.42 mmol/L (0.7–1.0)

eGFR >60 mL/min/1.73 m²(>60)

Which drug is most likely to be contributing to his symptoms?

- A.** Amlodipine
- B.** Bisoprolol
- C.** Eplerenone
- D.** Esomeprazole
- E.** Losartan



- 63.** A 42 year old woman has recent weight loss following an upper respiratory tract illness (URTI).

Her thyroid is slightly enlarged, smooth and diffusely tender.

Investigations:

Thyroid stimulating hormone 0.1 mU/L (0.3–4.2)

Free T4 35 pmol/L (9–25)

Free T3 11 pmol/L (4.0–7.2)

Thyroid technetium (Tc- 99m) pertechnetate scan shows generalised reduced uptake.

Which is the most likely explanation for these findings?

- A.** Graves disease
- B.** Non-specific response to stress of illness
- C.** Thyroid adenoma
- D.** Toxic thyroid nodule
- E.** Viral thyroiditis



- 64.** A 68 year old man with type 2 diabetes mellitus has a non-healing wound on his left foot following a minor injury 3 weeks ago.

His temperature is 38.3°C. The left foot is cool. There is a 2 cm ulcer on his toe, with cellulitis up to the forefoot. Foot pulses are palpable in both feet. He has reduced sensation to light touch in both feet.

Which is the most appropriate management?

- A.** Immediate admission to hospital
- B.** Refer to podiatrist
- C.** Refer to practice nurse for wound dressing
- D.** Start oral flucloxacillin
- E.** Urgent referral to diabetic outpatient clinic



65. A 35 year old woman has 2 months of palpitations and sweating.

A 2 cm nodule can be palpated in the left lobe of the thyroid gland, but the lymph nodes are not enlarged.

Investigations:

TSH <0.01 mU/L (0.3–4.2)

Free T4 29 pmol/L (9–25)

Free T3 8.5 pmol/L (4.0–7.2)

TSH-receptor antibody negative

Which investigation is most likely to establish the diagnosis?

- A.** CT scan of neck
- B.** Technetium uptake scan
- C.** Thyroid peroxidase antibody
- D.** Ultrasound-guided fine needle aspiration of nodule
- E.** Ultrasound scan of thyroid



66. A 45 year old man has had two episodes of headache, flushing and palpitation over the past 12 months. His BP has been noted to be very high during these episodes.

Investigations:

Plasma metanephrine 1200 pmol/L (<600)

Plasma normetanephrine 3500 pmol/L (<1000)

Which class of drug should be initiated as a first-line antihypertensive?

- A.** Alpha adrenergic receptor agonist
- B.** Alpha adrenergic receptor antagonist
- C.** Antimuscarinic agent
- D.** Beta adrenergic receptor agonist
- E.** Beta adrenergic receptor antagonist



- 67.** A 33 year old doctor who is training to be a surgeon develops an acute febrile illness accompanied by a generalised maculopapular rash.

Blood tests confirm acute HIV infection. He starts antiretroviral treatment. Occup Occupational health is overseeing his clearance and ongoing monitoring for surgical practice. He is concerned about the long-term impact on his surgical career.

Which advice is most appropriate regarding his future surgical practice?

- A.** He can perform exposure-prone procedures once adequate viral suppression is established
- B.** He can perform exposure-prone procedures once antiretroviral treatment is started
- C.** He can perform exposure-prone procedures provided each patient knows his diagnosis
- D.** He must permanently restrict his practice to procedures that are not exposure prone
- E.** He must transfer to a specialty that does not involve operative procedures



68. A 19 year old woman is found collapsed at home. She drinks one bottle of vodka a day. She is a person who injects drugs.

Her temperature is 35.2°C, pulse 125 bpm, BP 102/62 mmHg, respiratory rate 22 breaths per minute and oxygen saturation 96% breathing air. Her right leg is swollen and tender to touch.

She is catheterised and passes a small amount of dark reddish-brown urine. Urinalysis: blood 3+, protein 1+, nitrites negative.

Investigations:

Urea 23.1 mmol/L (2.5–7.8)

Creatinine 534 µmol/L (60–120)

Calcium (adjusted) 1.9 mmol/L (2.2–2.6)

Phosphate 3.6 mmol/L (0.8–1.5)

What is the most likely cause of her AKI?

- A.** Acute interstitial nephritis
- B.** ANCA-associated vasculitis
- C.** Pyelonephritis
- D.** Renal vein thrombosis
- E.** Rhabdomyolysis



69. A 37 year old woman has 5 hours of right-sided abdominal pain. The pain is sharp and radiates towards her groin. She is vomiting and is restless due to the pain. She is allergic to penicillin.

Investigations:

Urinalysis: blood 4+

β hCG: negative

Which is the most appropriate immediate management?

- A.** Co-amoxiclav
- B.** Diclofenac
- C.** Hyoscine butylbromide
- D.** Tamsulosin
- E.** Tramadol



70. A 55 year old man attends the Emergency Department with severe generalised headache of sudden onset.

He is in pain, photophobic and has a complete ptosis on the left side. On lifting the left eyelid, his eyeball is looking downwards and outwards, and his pupil is dilated and unresponsive to light.


Which is the most likely site of aneurysmal rupture?

- A.** Anterior communicating artery
- B.** Middle cerebral artery
- C.** Ophthalmic artery
- D.** Posterior cerebral artery
- E.** Posterior communicating artery



- 71.** A 42 year old man has blurred vision in his right eye, gradually worsening over 6 months. He has had no headache, diplopia, or systemic symptoms. His visual acuity is 6/9 right and 6/6 left.

His visual field shows a deficit in the right eye vision (see image). Fundi and remaining neurological examination are normal.



1493.18496

What is the most likely underlying cause?

- A.** Anterior ischaemic optic neuropathy
- B.** Chronic glaucoma
- C.** Internal carotid artery aneurysm
- D.** Pituitary macroadenoma
- E.** Temporal lobe glioma



72. A 22 year old woman has 2 days of worsening visual loss in her left eye and can now only detect hand movements. Six months earlier, she had an episode of facial numbness that resolved. She smokes 10 cigarettes per day and drinks three bottles of wine each week.

She has a relative afferent pupillary defect with visual acuity of hand movements in the left eye. Neurological examination is otherwise normal. Her BMI is 37 kg/m². MR scan of brain shows multiple T2 hyperintensities.

Which action is most likely to reduce her risk of a further neurological episode?

- A.** Alcohol reduction
- B.** IV methylprednisolone for 3 days
- C.** Oral methylprednisolone for 5 days
- D.** Smoking cessation
- E.** Weight loss



73. A 78 year old woman has had 6 months of progressive heaviness and aching in both legs when walking. Stopping and remaining upright provides little relief, but sitting down or leaning forwards relieves her symptoms. She can walk further when pushing a shopping trolley and finds walking uphill easier than downhill. She has hypertension and type 2 diabetes mellitus.

Her feet are warm with palpable pulses. Lower limb power and sensation are normal at rest. Lumbar extension reproduces her leg discomfort.

Which investigation is most likely to confirm the underlying diagnosis?

- A.** Ankle brachial pressure index
- B.** CT scan of lumbar spine
- C.** MR imaging of lumbar spine
- D.** Nerve conduction studies
- E.** X-ray of lumbar spine



- 74.** A 34 year old man has leg weakness, difficulty walking and back pain for 24 hours. He has tingling in both hands. He had a self-limiting diarrhoeal illness 10 days ago.

He has bilateral proximal leg weakness, with absent knee and plantar reflexes. Vibration sense is reduced to the hips and pin-prick to the knees bilaterally.

What is the most likely diagnosis?

- A.** Anterior spinal artery infarction
- B.** Guillain-Barré syndrome
- C.** Hypokalaemic paralysis
- D.** Spinal cord compression
- E.** Transverse myelitis



75. A 50 year old man has 4 weeks of persistent floaters in his right eye. Initially he could see flashing lights in this eye that have now resolved.

Retinal examination is normal on direct ophthalmoscopy.

Which is the most likely diagnosis?

- A.** Dry macular degeneration
- B.** Posterior vitreous detachment
- C.** Retinal detachment
- D.** Retinal tear
- E.** Vitreous haemorrhage



- 76.** A 31 year old man has had a chronic rash for several years which is gradually spreading (see image). Using moisturiser on his skin has made no difference.



Which is the most appropriate first line treatment?

- A.** Oral methotrexate
- B.** Oral psoralen with local ultraviolet A (PUVA) irradiation
- C.** Topical calcineurin inhibitor
- D.** Topical mild corticosteroid with antifungal agent
- E.** Topical potent corticosteroid plus topical vitamin D analogue



77. A 38 year old white woman has 6 months of facial flushing, red spots and pustules on her face.

She has telangiectasia over the cheeks with erythematous papules and pustules.

Which is the most likely diagnosis?

- A.** Acne vulgaris
- B.** Discoid lupus erythematosus
- C.** Rosacea
- D.** Sarcoid
- E.** Seborrhoeic dermatitis



- 78.** A 67 year old patient has fatigue for a few months. There is no change in weight and no pain. They are not on any regular medication or over-the-counter drugs.

Investigations:

Haemoglobin 145 g/L (130–175)

Erythrocyte sedimentation rate 4 mm/h (<20)

Creatinine 78 $\mu\text{mol/L}$ (60–120)

Calcium 2.8 mmol/L (2.2–2.6)

Vitamin D 81 nmol/L (75–100)

TSH 3.5 mU/L (0.3–4.2)

Parathyroid hormone 17.1 pmol/L (1.6–8.5)

24 h urine calcium excretion normal

Which is the most likely diagnosis?

- A.** Familial hypocalciuric hypercalcemia
- B.** Multiple myeloma
- C.** Primary hyperparathyroidism
- D.** Secondary hyperparathyroidism
- E.** Tertiary hyperparathyroidism



79. A 70 year old woman has 1 month of worsening memory and mild confusion. She has a history of hypertension. She takes lisinopril and atorvastatin. She smokes 20 cigarettes per day.

She is disorientated in time and place. She has weakness of hip flexion in both legs. Her Montreal Cognitive Assessment score is 22/30, with deficits in recall and attention.

Investigations:

MR scan of brain: T2 weighted image shows several hyperintense lesions at border of grey and white matter in frontal and temporal lobes

Which is the most likely diagnosis?

- A.** Brain metastases
- B.** Frontotemporal dementia
- C.** Glioblastoma multiforme
- D.** Progressive multifocal leucoencephalopathy
- E.** Vascular dementia



- 80.** A 25 year old man has gradually worsening pain and swelling of his left testicle over 2 days. He also reports dysuria and a small amount of urethral discharge. He had unprotected sexual intercourse with a new partner 2 weeks ago.

His temperature is 37.8°C and pulse 88 bpm. There is tenderness and swelling behind his left testicle, extending into the testicle itself. The testicle has a normal lie.

Which is the most likely diagnosis?

- A.** Epididymo-orchitis
- B.** Hydrocele
- C.** Inguinal hernia
- D.** Testicular cancer
- E.** Testicular torsion



- 81.** A 15 year old boy has acute onset of left testicular pain and lower abdominal pain associated with nausea and vomiting over the past 90 minutes. There is no dysuria or urethral discharge.

His temperature is 37.6°C, pulse 86 bpm, BP 150/93 mmHg, respiratory rate 14 breaths per minute and oxygen saturation 98% breathing air. There is left hemiscrotal erythema, and the left testicle is swollen, tender and high riding with a transverse lie. Urinalysis is normal.

Which is the most likely diagnosis?

- A.** Epididymitis
- B.** Inguinal hernia
- C.** Orchitis
- D.** Testicular tumour
- E.** Torsion of testis



- 82.** A 28 year old woman has a painful, red, watery left eye. She wears contact lenses but has been unable to wear them for several days because of the pain. Over the past 24 hours, she has been shying away from bright light.

She has localised erythema surrounding the medial aspect of the cornea of her left eye. There is dendritic fluorescein staining under UV light over her medial cornea.

Which is the most likely diagnosis?

- A.** Adenoviral conjunctivitis
- B.** Anterior uveitis
- C.** Bacterial conjunctivitis
- D.** Herpetic keratitis
- E.** Traumatic corneal ulceration



- 83.** A 36 year old man has lost weight and experienced night sweats and a dry cough over 6 weeks.

His temperature is 37.3°C. He has a palpable right cervical lymph node. His chest is clear.

Investigations:

Chest X-ray: opacification in right upper lobe

Lymph node biopsy: lymph node architecture is effaced by an infiltrate composed of macrophages with strong epithelioid features with central caseous necrosis

Which is the most likely diagnosis?

- A.** Histoplasmosis
- B.** Lymphoma
- C.** Sarcoidosis
- D.** Seminoma
- E.** Tuberculosis



84. A 32 year old woman has had 2 days of difficulty chewing and cannot finish a whole meal. She has also had 6 weeks of double vision. This has been progressively worsening and is most noticeable at the end of the day.

What is the most likely diagnosis?

- A.** Botulism
- B.** Graves' disease
- C.** Motor neurone disease
- D.** Myasthenia gravis
- E.** Oculopharyngeal muscular dystrophy



85. A 25 year old woman has 4 months of dull frontal headaches that are increasing in frequency. She now has symptoms on most days. Her headache is worse in the morning. She has associated nausea, fleeting loss of vision and tinnitus. She takes paracetamol as required, with some relief. Her BMI is 32 kg/m².

Which is the most likely diagnosis?

- A.** Cerebral venous sinus thrombosis
- B.** Idiopathic intracranial hypertension
- C.** Medication overuse headache
- D.** Migraine
- E.** Tension headache



86. A 58 year old man has one episode of painless visible haematuria. He has hypertension and takes amlodipine. He smokes 10 cigarettes per day.

His pulse is 72 bpm and BP 140/92 mmHg. Abdominal examination is normal. Urinalysis: blood 2+.

Investigations:

Haemoglobin 149 g/L (130–175)

Creatinine 105 $\mu\text{mol/L}$ (60–120)

eGFR 63 mL/min/1.73 m²(>60)

Mid-stream urine: no growth

What is the most appropriate next step in management?

- A.** Cystoscopy
- B.** Repeat urine culture in 1 month
- C.** Urine cytology
- D.** Urine protein:creatinine ratio
- E.** US scan of abdomen



- 87.** A 62 year old man has difficulty walking because of increasing leg swelling. He smokes 20 cigarettes daily.

His BP is 144/94 mmHg, and he has oedema of both legs with swelling of his scrotum. Urinalysis shows blood 1+, protein 4+.

Investigations:

Urea 16.0 mmol/L (2.5–7.8)

Creatinine 180 μ mol/L (60–120)

Urine protein: creatinine ratio 680 mg/mmol (<30)

Albumin 22 g/L (35–50)

What investigation is most likely to establish the underlying diagnosis?

- A.** 24 hour urine collection
- B.** CT abdomen and pelvis
- C.** Cystoscopy
- D.** Renal biopsy
- E.** Serum anti-neutrophil cytoplasmic antibodies



- 88.** A 54 year old woman has had worsening headaches for 4 months. Over the past 3 years, her rings have become tight and her shoe size has increased. She also reports joint pain and excessive sweating.

Her serum insulin-like growth factor 1 (IGF-1) is mildly above the age-adjusted reference range on two measurements.

Which is the most appropriate next diagnostic investigation?

- A.** Insulin tolerance test
- B.** Mixed meal test
- C.** Oral glucose tolerance test with growth hormone measurements
- D.** Prolonged supervised fast
- E.** Short Synacthen® test



- 89.** A 20 year old man has retinitis pigmentosa. His parents are unaffected, but his maternal grandfather had retinitis pigmentosa as did the maternal grandfather's brother.

Which is the most likely pattern of inheritance?

- A.** Autosomal recessive
- B.** Multifactorial
- C.** Sporadic mutation
- D.** X-linked dominant
- E.** X-linked recessive



90. A 74 year old woman is admitted with fever, dysuria and new confusion. She has no drug allergies.

Her temperature is 38.2°C, pulse 110 bpm and BP 90/60 mmHg. Blood and urine cultures are taken. Intravenous fluids and ceftriaxone are started.

The following day, blood and urine cultures both grow *Enterococcus faecalis*. Antibiotic susceptibility results are pending.

Which antibiotic should replace ceftriaxone while susceptibility results are pending?

- A.** Intravenous amoxicillin
- B.** Intravenous ciprofloxacin
- C.** Intravenous metronidazole
- D.** Oral doxycycline
- E.** Oral nitrofurantoin



91. A 66 year old man has a 1 year history of increasing urinary hesitancy, a weak stream and terminal dribbling. He wakes three to four times each night to pass urine. He has no dysuria, visible haematuria or previous urinary retention.

Digital rectal examination reveals a smoothly enlarged, non-tender prostate without nodules.

Investigations:

Urinalysis: normal

Urine culture: no growth

Creatinine 80 $\mu\text{mol/L}$ (60–120)

Which additional investigation is most appropriate to offer as part of his initial assessment?

- A.** Cystoscopy
- B.** Prostate biopsy
- C.** Prostate-specific antigen
- D.** Ultrasound scan of prostate
- E.** Urodynamic studies



92. A 25 year old woman is 16 weeks pregnant. She has a new kitten and has been advised not to handle cat litter.

What infection is a risk to the fetus?

- A.** Cytomegalovirus
- B.** Group B streptococcus
- C.** Hepatitis B
- D.** Toxoplasmosis
- E.** Varicella zoster virus



- 93.** An 83 year old man on a rehabilitation unit reports urinary hesitancy, a weak stream and prolonged terminal dribbling that wets his clothing. These symptoms have gradually worsened over 6 months. He has no urgency, dysuria or suprapubic pain.

His bladder is not palpable. Digital rectal examination reveals a smoothly enlarged prostate. His urinalysis is normal.

Investigations:

Creatinine 86 $\mu\text{mol/L}$ (60–120)

Post-void residual urinary volume: 130 mL

His symptoms remain troublesome despite advice on fluid timing and double voiding.

Which is the most appropriate initial drug treatment?

- A.** Alfuzosin
- B.** Finasteride
- C.** Furosemide
- D.** Mirabegron
- E.** Tolterodine



- 94.** A 52 year old woman with type 2 diabetes mellitus has had burning and tingling pain in both feet for 6 months. The pain is worse at night and disrupts her sleep. She has ischaemic heart disease and atrial fibrillation.

There is symmetrical loss of pinprick and vibration sensation in a stocking distribution. Her feet are warm, with palpable pulses and no ulcers.

Investigations:

HbA1c 87 mmol/mol (20–42)

Estimated GFR 78 mL/min/1.73 m² (>60)

Which medication is most appropriate to start for her pain?

- A.** Amitriptyline
- B.** Gabapentin
- C.** Ibuprofen
- D.** Morphine
- E.** Prednisolone



95. A 19 year old man presents to the Emergency Department with 48 hours of fever, sore throat and a generalised rash that began on his face and spread downwards.

His temperature is 39.1°C, pulse rate 112 bpm and BP 106/65 mmHg. He has a widespread blanching rash and conjunctivitis. He has white spots in his mouth.

Which is the most likely diagnosis?

- A.** Erythrovirus (parvovirus) B19
- B.** Measles
- C.** Meningococcal septicaemia
- D.** Rubella
- E.** Scarlet fever



96. A 19 year old woman has 2 days of malaise and headache. She has recently returned from a 2 week holiday to India.

Her temperature is 38°C and pulse rate 77 bpm. She has photophobia and neck stiffness.

Investigations:

Cerebrospinal fluid: Opening pressure 90 mmH₂O (50–180)

Total protein 1.2 g/L (0.15–0.45)

Glucose 4.1 mmol/L (2.2–4.4)

Cell count 500 /μL (<5)

Lymphocyte count 90% (60–70)

Paired plasma glucose 6.0 mmol/L

Which is the most likely diagnosis?

- A.** Bacterial meningitis
- B.** Cryptococcal meningitis
- C.** Malaria
- D.** Tuberculous meningitis
- E.** Viral meningitis



97. A 65 year old woman has had polyuria and polydipsia for 3 months. She has a history of bipolar disorder, hypertension and peptic ulcer disease. She takes lisinopril, lithium and omeprazole. She is a non-smoker and drinks around 30 units of alcohol per week.

Investigations:

Sodium 148 mmol/L (135–146)

Potassium 3.9 mmol/L (3.5–5.3)

Urea 10.1 mmol/L (2.5–7.8)

Creatinine 79 μ mol/L (60–120)

Serum osmolality 308 mOsmol/kg (285–295)

Calcium 2.3 mmol/L (2.2–2.6)

HbA1c 38 mmol/mol (20–42)

Urinary osmolality 100 mOsmol/kg (100–1000)

Urinary sodium 65 mmol/L (>20)

She is given a therapeutic trial of oral desmopressin, and the investigations above remain unchanged.

Which is the most likely diagnosis?

- A.** Arginine vasopressin deficiency (AVPD, cranial diabetes insipidus)
- B.** Arginine vasopressin resistance (AVPR, nephrogenic diabetes insipidus)
- C.** Primary polydipsia
- D.** Syndrome of inappropriate ADH secretion
- E.** Type 2 diabetes mellitus



98. A 52 year old man has been generally unwell with pain in the right eye for 5 days.

He cannot open his right eyelid, and the right pupil is dilated and does not react to light. His right eye is deviated downwards and laterally, and he cannot move the eye upwards or medially. On attempting to look down further, his right eyeball rotates slightly.

Which is the best description of the clinical signs?

- A.** Complete third nerve palsy
- B.** Complete third nerve palsy and fourth nerve palsy
- C.** Fourth nerve palsy
- D.** Horner's syndrome
- E.** Partial third nerve palsy



99. A 35 year old man has sudden loss of vision in his right eye with no prodromal symptoms. He has longstanding type 1 diabetes and has not attended for eye screening for several years.

Which complication of diabetes is the most likely cause of his visual symptoms?

- A.** Cataract
- B.** Ischaemic optic neuropathy
- C.** Maculopathy
- D.** Retinal detachment
- E.** Vitreous haemorrhage



100. A 32 year old man has had persistent non-visible haematuria for 3 months, initially detected during an army medical assessment. He has had no visible haematuria. He had no family history of renal disease. He drinks 30 units of alcohol a week.

His pulse is 74 bpm and BP 147/93 mmHg. Examination is normal. Urinalysis shows blood 3+ and protein 2+ only.

Investigations:

Creatinine 104 $\mu\text{mol/L}$ (60–120)

eGFR $>60 \text{ mL/min/1.73 m}^2(>60)$

ANA negative

US scan of kidneys: normal appearance

Which is the most likely diagnosis?

- A. Alport's syndrome
- B. IgA nephropathy
- C. Lupus nephritis
- D. Membranous nephropathy
- E. Thin basement membrane disease